

NAME.....INDEX NO.....
CANDIDATE'S SIGNATURE.....DATE.....
SCHOOL.....

233/3
CHEMISTRY
PAPER 3
PRACTICALS
MAY/JUNE 2014
TIME: 2 ¼ HOURS

EKSIKA JOINT EVALUATION TEST.

Kenya Certificate of Secondary Education (K.C.S.E)

233/3
CHEMISTRY
PAPER 3
PRACTICALS
MAY/JUNE 2014
TIME: 1 ¼ HOURS

INSTRUCTIONS TO CANDIDATES.

- Write your name and index number in the spaces provided above.
- Sign and write the date of examination in the spaces provided above.
- Answer **ALL** questions in the spaces provided above.
- All workings **MUST** be clearly shown where necessary.
- You are not allowed to work with the apparatus for the first 15 minutes of the 2 ¼ Hours allowed for this paper. This time is to enable you read the question paper and make sure you have all the chemicals and the apparatus that you may need.
- Mathematical tables and silent electronic calculators may be used.

FOR EXAMINERS' USE ONLY.

Question	Maximum Score	Candidates' Score
1	12	
2	40	
3	14	
	40	

*This paper consists of 4 printed pages.
Candidates should check the questions paper to ascertain that all pages are printed as indicated
and no questions are missing.*

- 1 You are provided with:
 Solution M 0.2M hydrochloric acid,
 Solution F containing 15.3g per litre of basic compound $G_2X.H_2O$.
 You are required to determine the relative atomic mass of G.

PRECEDURE:

Place solution M in a burette ,pipette 25cm^3 of solution F into a 250cm^3 conical flask. Add two drops of methyl orange indicator and titrate. Record your results in the table below.
 Repeat the procedure two more times and complete table I.

Table I

a) i)

	I	II	III
Final burette reading			
Initial burette reading			
Volume of solution M used (cm^3)			

(4mks)

ii) What is the average volume of solution M.?

(1mk)

.....

b) Given that one mole of F reacts with 2moles of M. Calculate the;

i) number of moles the basic compound, $G_2X, 10H_2O$ in the volume of solution F used. (2mks)

.....

ii) Concentration of solution F in mole per litre.

(2mks)

.....

iii) Relative formula mass of the basic compound, $G_2X.10H_2O$.

(1 ½ mks)

.....

- iv) relative atomic mass of G (Relative formula Mass of $X=60$, atomic mass of $H=1.0$, $O=16.0$). (1 ½ mks)

.....
.....
.....

2 You are provided with:

- 1 1.899g of solid P, solid P is adiabatic acid H_2X .
- 2 0.5M Solution of the dibasic acid , H_2X , Solution V.
- 3 Sodium hydroxide, Solution K.

You are required to determine:

- a)
 - i) the molar heat of solid P.
 - ii) the heat of reaction of one mole of the dibasic acid with sodium hydroxide.
- b) Calculate the heat of reaction of solid H_2X with aqueous sodium hydroxide.

PROCEDURE I.

Place 30cm^3 of distilled water into a 100ml beaker. Measure the initial temperature of the water and record it in the table II below. Add all the solid P at once; stir the mixture carefully with the thermometer until all the solid dissolves. Measure the final temperature reached and records it in the table II

Table II

Final temperature ($^{\circ}\text{C}$)	
Initial temperature ($^{\circ}\text{C}$)	

- a) Determine the change in temperature ΔT_1 (1½mks)

.....

b) Calculate the:

- i) heat change when H_2X dissolves in water, (Assuming the heat capacity of the solution is $4.2\text{Jg}^{-1}\text{K}^{-1}$ and density is 1g/cm^3) (2mks)
- ii) number of moles of the acid that were used. (Relative formula mass of H_2X is 126) (1mk)

.....
.....
.....

- iii) molar heat of solution ΔH_1 solution of the acid H_2X . (1mk)

.....
.....
.....
PROCEDURE II.

Place 30cm³ of solution V into a 100cm³ beaker. Measure the initial temperature and record it in table III below. Measure 30cm³ of sodium hydroxide, solution K. Add all of the 30cm³ of solution K at once to V in the beaker. Stir the mixture with the thermometer. Measure the final temperature reached and record it in table III.

Table III.

a)

Final temperature (°C)	
Initial temperature (°C)	

(1 ½ mks)

b) Determine the change in temperature, ΔT_2 . (½ mk)


c) Determine the:

i) heat change for the reaction (Assume the heat capacity of the solution is 4.2Jg⁻¹k⁻¹ and density is 1g/cm³) (2mks)

ii) Number of moles of the acid used (H₂X). (1mk)

iii) Heat of reaction , ΔH_2 of one mole of the acid H₂X with sodium hydroxide (1mk)

d) Given that,



Calculate ΔH_3 for the reaction $H_2X_{(s)} + 2OH^{-1}_{(aq)} \longrightarrow 2H_2O_{(l)} + X^{2-}_{(aq)}$ (2mks)

.....

3 You are provided with solid S. Carry out the tests below and record your observations and inferences in the spaces provided.

a) Place about one third of solid S in a dry test tube. Heat the solid gently and the strongly. Test any gases produced with blue and red litmus papers.

Observations	Inferences
(2mks)	(1mk)

b) Dissolve the remaining portion of solid S in $8cm^3$ of distilled water.

i) Divide the solution into the first portions, to the first portion, add aqueous sodium hydroxide drop wise until in excess.

Observations	Inferences
(1mk)	(2mks)

ii) To the second portion , add aqueous ammonia dropwise in excess.

Observations	Inferences
(1mk)	(1mk)

iii) To the third portion , add 10cm^3 of barium chloride solution.

Observations	Inferences
(1mk)	(1mk)

iv) To the forth portion , add about 1cm^3 of Lead (II) nitrate solution.

Observations	Inferences
(1mk)	(1mk)

v) To the fifth portion, add about 2ml of hydrogen peroxide then about 1cm^3 of sodium hydroxide solution.

Observations	Inferences
(1mk)	(1mk)

ANSWERS:

Order a copy of answers from www.schoolsnetkenya.com/order-e-copy

NB> We charge Kshs. 100 ONLY to meet website, e-resource compilation and provision costs