

---

**KENYA NATIONAL EXAMINATION COUNCIL**  
**REVISION MOCK EXAMS 2016**  
**TOP NATIONAL SCHOOLS**

**ALLIANCE GIRLS HIGH SCHOOL**

**232/1**

**PHYSICS**

**PAPER 3**

**SCHOOLS NET KENYA**

Osiligi House, Opposite KCB, Ground Floor

Off Magadi Road, Ongata Rongai | Tel: 0711 88 22 27

E-mail: [infosnkenya@gmail.com](mailto:infosnkenya@gmail.com) | Website: [www.schoolsnetkenya.com](http://www.schoolsnetkenya.com)

**ALLIANCE GIRLS HIGH SCHOOL KCSE TRIAL  
AND PRACTICE EXAM 2016  
Paper 3 (Practical)**


---

**QUESTION 1 PART A**

1. You are provided with the following
- A micrometer screw gauge (to be shared)
  - A voltmeter (0-3v or 0-5v)
  - Ammeter (0-1A)
  - A switch
  - A jockey/long wire with crocodile clip attached
  - One new dry cell
  - 8 connecting wires with crocodile clips attached to one end
- Proceed as follows*

(a) Set up the circuit below, Fig 1 ensure that when the switch is open, both meters read zero, keep the switch open when readings are not being taken

**Fig 1**



- (i) Measure and record the diameter  $d$  of the nichrome wire AB using the micrometer screw gauge  
 $d = \text{m}$  ( ½ mk)
- (ii) Disconnect the jockey from wire AB and close the switch. Record the value  $E$  of the voltmeter reading.  
 $E = \text{V}$  ( ½ mks)
- (b) Now, connect the jockey on AB at a distance  $L = 2.5\text{cm}$ . Close the switch and record the voltmeter and ammeter readings,  $V$  and  $I$  respectively in table 1 below.




Table 1

L(cm)	2.5	7.5	10.0	20.0	30.0	40.0
D.d(v)						
Current I(A)						
IV(watts)						

(i) Complete the table (5mks)

(ii) Plot a graph of IV (vertical axis) against L (5mks)

(iii) Using your graph, find the value  $l_0$  where your graph (the horizontal axis)

$l_0 = \underline{\hspace{2cm}}$  cm (1mk)

(c)(i) Now, place the jockey on AB such that long the L is equal to the value of  $l = 63$ cm. close the switch and record both the voltmeter reading, V and the ammeter reading, I

$V = \underline{\hspace{2cm}}$  V (½ mk)

$I = \underline{\hspace{2cm}}$  A (½ mk)

(ii) Work out the values v where (1mk)

$$r = \frac{E - V}{I}$$

(d) Work out the value of e where (1 ½ mks)


$$e = \frac{\pi r d^2}{2.52}$$

**Question 1 part B**

You are provide with

- Rectangular glass block
- Four optical pins
- Ruler
- Soft board
- Plain paper
- Cellotape
- Vernier calipers (to be shared)

(e) Set up the apparatus as shown in figure below



Proceed as follows

- (f) (i) Using the vernier calipers provided, measure and record the breadth  $b$  of the glass block  
 $b = \underline{\hspace{2cm}}$  cm (½ mk)
- (ii) Using cellotape, fix the mirror on one side (length) of the glass block and trace its outline on the plain paper
- (iii) Draw the normal  $NK$  to the side  $AB$  and measure angle  $i = 10^\circ$  from the normal
- (iv) Draw the line representing the incident ray and fix pins  $P_1$  and  $P_2$  as shown in the figure
- (v) By observing the images of the pins  $P_1$  and  $P_2$ , locate the position  $P_3$  and  $P_4$  such that they appear in a line (no parallax) using other pins
- (vi) Join the points  $P_3$  and  $P_4$  and extend them to intersect line  $P_1P_2$  produced. Measure the perpendicular distance  $Y$
- (vii) Repeat steps (iii-vi) for different values of  $i$  given and record your values in the table 2 below

Table 2

$i^\circ$	10	20	30	40
$Y(\text{cm})$				

- (g) (i) Determine the average of the values of  $Y$  (1mk)
- (ii) Determine the values of constant  $k$  given that  $k = \frac{b}{Y}$  (1mk)

## 2. PART A

You are provided with the following

- Metre rule
- Knife edge
- 10 microscope slides
- A 50g mass
- A piece of cellotape
- A pair of vernier calipers

Proceed as follows

- (a) Using the vernier calipers provided measure the length  $l$  and the width of the microscope slide

L= \_\_\_\_\_  
 W= \_\_\_\_\_ (1mk)

(b) Stack ten(10) slides together using a cello tape as shown below fig 3




Figure 3

(i) Measure the thickness T of the space of microscope slab (1mk)


T= \_\_\_\_\_

(ii) Determine the volume v of the stack (1mk)

V=(WT)= \_\_\_\_\_

(c) Balance the metre rule at its centre of gravity and maintain the position of the fulcrum on the centre of gravity throughout the experiment

Place the 50g mass and the stack of slides as shown in figure 4 below



Adjust the position of both the spaces and the mass until the rule is again balanced make the distances x and y as large as possible

(i) x= \_\_\_\_\_ (1mk)

y= \_\_\_\_\_

(ii) Calculate the mass M in grams of the stick of slides given that (1mk)

$$m = 50 \frac{x}{y}$$


(iii) Determine the density of glass given that  $density = \frac{m}{v}$  (1mk)

**PART B**

(d) You are provided with a metre rule, a lens holder, convex lens, a candle amounted white screen proceed as follows

(e) Set up the apparatus as shown in figure 5 below

(f) Ensuring that L=100cm adjust the lens until you get a sharp diminished image on the screen. Measure the object distance u, and image distance v



(g)(i) Repeat the procedure with L=95cm, 90cm, 85cm,80cm and 75cm each time recording the value of u and v and tabulating the results in the table II below (5mks)

L(cm)	100	95	90	85	80	75
U(cm)						

V(cm)						
$m = \frac{v}{u}$						

(ii) Plot a graph of m against v

(5mks)

(iii) Determine the slope of the graph

(2mks)

(iv) Given that  $\frac{v}{f} = m + 1$ , determine the focal length of the lens from the graph above

(2mks)