
KENYA NATIONAL EXAMINATION COUNCIL
REVISION MOCK EXAMS 2016
TOP NATIONAL SCHOOLS

KABARAK HIGH SCHOOL
CHEMISTRY
PAPER 1
MARKING SCHEME


SCHOOLS NET KENYA
Osiligi House, Opposite KCB, Ground Floor
Off Magadi Road, Ongata Rongai | Tel: 0711 88 22 27
E-mail: infosnkenya@gmail.com | Website: www.schoolsnetkenya.com

KABARAK HIGH SCHOOL KCSE TRIAL AND PRACTICE EXAM 2016

QUESTION PAPER 1

MARKING SCHEME

- (a) Atoms of the same element with same atomic number but different mass no.s/neutrons
(b) Total sum of protons and neutrons
(c) Nuclides with same mass no.s but different atomic no.s
- (a) Calcium oxide reacts with chlorine to form calcium hypochlorite to form calcium hypochlorite
(b) Anhydrous CaCl_2 / conc. H_2SO_4
- (a) Both ammonia and water are polar molecules and hydrogen bonds are formed.
(b) Co-ordinate bond
- (a)(i) Naphthalene particles absorb heat increasing the kinetic energy increasing the vibrations.
(ii) Heat absorbed is used to break the bonds holding the solid particles together.
- (a) Enthalpy change when one mole of the compound is formed from its constituent elements in their standard states (stp)
(b) Molar enthalpy change of formation
Molar enthalpy change of combustion




$$\Delta H_1 + \Delta H_2 = \Delta H_3 \quad (\frac{1}{2})$$

$$\Delta H_1 + \Delta H_2 - \Delta H_2 = \Delta H_3 - \Delta H_2 \quad (\frac{1}{2})$$


$$\Delta H_1 = 3(-394) + 4(-286) - (-1202) \quad (\frac{1}{2}) = -124 \text{ kJ mol}^{-1} \quad (\frac{1}{2})$$

- (a) Pipe nearest ground level
- I
(b) (i) Vanadium (v) oxide/platinum
(iii) I yield decreases extra heat decomposes SO_3 /forward rxn is exothermic/equilibrium shifts to the left.
II Yields increases since extra oxygen is used
- In diamond all the bonds are strong covalent bonds while graphite structure of layers that held together by weak van der Waals forces that are easily broken

8. (a)



(b)



9. (i) I 1 and III
II II and IV

(ii) IV, has a double bond

10. (a) KBr

(b) $60 - 55 = 5\text{g}$

(c) Fractional crystallization

11. (a) - Thistle funnel to touch reagents
- Delivery tube to go through the hole of beehive shelf

(b) Sodium peroxide

12. (i) Bitumen, last to be collected;

(ii) Fractional distillation

(iii) Limited supply of oxygen

13. (a) $\text{H}^+_{(\text{aq})} + \text{OH}^-_{(\text{aq})} \longrightarrow \text{H}_2\text{O}_{(\text{l})}$




(b) I Y_2 – complete neutralization /end point

II Y_1 and Y_2 –Neutralization is taking place producing heat.

III Y_2 and Y_3 reaction has come and products are cooling/cooling releases heat to the surrounding.


Heat is given out	
Purple vapour condenses on cooler parts of test tube into shiny grey crystals	Physical
Solid decomposes to form gas and black solid/blue-green due to loss of water	Chemical

- 15 (a) A base that dissolves in water to give hydroxide ions.
 (b)(i) Very soluble in water, thus it would dissolve in water instead of being collected.
 (ii) Less dense than air
- 16 (a) Conc. Sulphuric acid rejected if concentrated missing
 (b) $\text{H}_2\text{SO}_4(l) + \text{NaNO}_3(s) \longrightarrow \text{NaHSO}_4(s) + \text{HNO}_3(g)$
 (c) Prevent decomposition of nitric acid
- 17 (a)(i) X-Oxygen
 Y-Hydrogen
 (b) Water is a weak electrolyte while hydrogen chloride gas in water is a strong electrolyte
 Water has no ions but $\text{HCl}(aq)$ has ions
- 18 (a) K and N; same group/same valence electrons/lose two electrons
- 19 - copper
 - the two electrodes have the same electrode potentials
 (b) A
 Has the most negative reduction potential, so its tendency to donate electrons is the highest.
- (c) $\text{B}(s) \longrightarrow \text{B}^{2+}(aq) + 2e^-$
- 20 (a) At constant temperature, the volume is inversely proportional to the pressure.
 (b) $P_1V_1 = P_2V_2$ ($\frac{1}{2}$)
 $3 \times 1 = 2 \times V_2$ ($\frac{1}{2}$)
 $V_2 = \frac{3 \times 1}{2} = 1.5$ (1)
- 21 W- mixture of sodalime and sodium ethanoate
 P- Methane
 (b) Substitution
22. (a) Time taken for half the amount to decay
 (b) $100 \xrightarrow{t_{1/2}} 50 \xrightarrow{t_{1/2}} 25 \xrightarrow{t_{1/2}} 12.5 \longrightarrow$
 $3 t_{1/2} = 15.6 \text{ yrs}$ (1)
 $t_{1/2} = 5.2 \text{ yrs}$ (1)
- 23 (a) charge that atoms have in molecules or ions
 (b) $\text{Cr}_2\text{O}_7^{2-}$ $2\text{Cr} + 7(-2) = -2$
 $2\text{Cr} = +12$
 $\text{Cr} = +6$
 (b) MnO_4^- $\text{Mn} + 4(-2) = -1$
 $\text{Mn} = +7$
- 24 (a)(i)
 H
 |



(c) Methyl Methanoate

(d) Few drops of concentrated H_2SO_4



$$Q = 0.5 \times 18 \times 60 \times \frac{1}{2}$$

$$= 540 \text{ c } \sqrt{\frac{1}{2}}$$

$$\text{if } \longrightarrow 108\text{g}$$

$$96500\text{c} \longrightarrow \frac{108}{96500} \sqrt{\frac{1}{2}}$$

$$540\text{c} \longrightarrow \frac{108}{96500} \times 540$$

$$= 0.6044\text{g}$$