CELL PHYSIOLOGY - ANSWERS

- 1. a) i) Diffusion
 - ii) Active transport
 - b) Diffusion-A concentration gradient between sodium ions in sap and those in the pond.

Active transport-energy in form of ATP must be available/Oxygen and food in the living tissue for respiration provide energy.

2. A film of water surrounds the soil particle. Root hairs of the plants

penetrate between the soils particles/are close to the soil particles; cell sap of the root hair cells is more concentrated in solutes/has less water than the soil solution. Thus water moves into root hair cell by osmosis i.e across the cell a wall and the semi permeable membrane.

- 3. The leaves expose a smaller surface area to the sun. Thus reducing transpiration/excessive water loss.
- 4 a) Diffusion is defined as the net movement of a substance from a region where its concentration is high to a region where its concentration is low.
 - b) i) Diffusion gradient-the greater the diffusion gradient, the greater the rate of diffusion
 - ii) Surface area to volume ratio-the greater the S.A.V.R thehigher the temperature the greater the rate of diffusion.
 - iii) Temperature –The higher the temperature the greater the rate of diffusion
 - c) i) Absorption of mineral salts from the soil by root hairs
 - ii) Re-absorption of glucose molecules in the kidney tubule.

- iii) Absorption of digested food in the ileum e.g glucose, amino acids.
- 5. i) Uptake of water from the soil into root hairs of plant roots
 - ii) Movement of water from the veins of leaves through the leaf cellsto the atmosphere during transpiration.
- a) The visking tubing was fully filled with solution. Level of water in beaker decreased .
- b) Sucrose solution in visking tubing created high concentration gradient.

-Water molecules moved from distilled water to the visking tubing by osmosis.

Plant cells have cells membrane and cell wall. When the cell is placed or
immersed in distilled water, the water is absorbed by osmosis. As cell becomes turgid,
the cell created an inward force, wall pressure that prevents the cell from bursting.

8.

Diffusion	Osmosis
Involves movement of particles of	Involves movement of solvent
molecules of liquid or gas.	It takes place through a semi-
It may be through a membrane or	permeable.
in air.	Rate affected by pH changes.

Compiled and supplied online by Schools Net Kenya | P.O. Box 85726 – 00200, Nairobi Tel:+254202319748 | +254 733 836593 | email: <u>infosnkenya@gmail.com</u> Order answers online at: <u>www.schoolsnetkenya.com</u>

- 9. a) Isotonic solution- a solution which has the same concentration as the cell sap.
 - b) Hypotonic solution- a solution which is less concentrated than the cell sap.
 - c) Hypertonic solution- A solution which is more concentrated than the cell sap.

10. Plants normally grow in soils whose solute concentration is lower than that of the cell sap. This enables the plants to take up water by osmosis. Addition of large amounts of salt to the soil increases the solute concentration of soil water beyond that of the cell sap. The result is that the plants lose water to the soil by osmosis. Since water is very important for maintaining the structural and metabolic activities of plants, its deficiency leads to death of the plants.

- a) The red blood cells take in water by osmosis. They swell and
 exert pressure on the fragile plasma membrane which then breaks. Plant cells
 take in water and swell but do not burst. This is because their tough cell wall can
 only stretch to a limited extent. Once fully stretched, the cell wall resists further
 expansion of the cell and no more water is taken up.
 - b) Fresh water protozoa take in water by osmosis. The excess wateris then actively pumped into the contractile vacuole which discharges the waterto the outside.